

Lefschetz-thimble path integral for studying the Silver Blaze phenomenon

Yuya Tanizaki

Department of Physics, The University of Tokyo

Theoretical Research Division, Nishina Center, RIKEN

Nov 25, 2015 @ KEK

Collaborators: Yoshimasa Hidaka, Tomoya Hayata (RIKEN)

Introduction and Motivation

Where is the finite-density QCD?

Neutron star

- Cold and dense nuclear matters
- $2m_{\text{sun}}$ neutron star (2010)
- Gravitational-wave observations (2017~)

Heavy-ion collision

- Low-energy scan of heavy-ion collisions are planned and run in many facilities (RHIC, SIS, J-PARC)

Reliable theoretical approach to **equation of state** must be developed!


$$Z(T, \mu) = \int \mathcal{D}A \text{Det}(\not{D}(A) + m) \exp -S_{\text{YM}}(A).$$

Silver Blaze problem in finite-density QCD

QCD & $|\text{QCD}|$

$$Z_{\text{QCD}} = \int \mathcal{D}A (\det \gamma_\nu D_\nu) e^{-S_{\text{YM}}}, \quad Z_{|\text{QCD}|} = \int \mathcal{D}A |\det \gamma_\nu D_\nu| e^{-S_{\text{YM}}}.$$

At $\mu = 0$, these two are the same! But,



At $T = 0$, the state must be equal to the QCD vac. for $|\mu| \lesssim m_N/3$.
Can we describe this “trivial” behavior using path integral?

Lefschetz decomposition formula

Oscillatory integrals with **many variables** can be evaluated using the “steepest descent” cycles \mathcal{J}_σ : (classical eom $S'(z_\sigma) = 0$)

$$\int_{\mathbb{R}^n} d^n x \, e^{-S(x)} = \sum_{\sigma} \langle \mathcal{K}_\sigma, \mathbb{R} \rangle \int_{\mathcal{J}_\sigma} d^n z \, e^{-S(z)}.$$

\mathcal{J}_σ are called Lefschetz thimbles, and $\text{Im}[S]$ is constant on it:

$$\mathcal{J}_\sigma = \left\{ z(0) \left| \lim_{t \rightarrow -\infty} z(t) = z_\sigma \right. \right\}, \quad \frac{dz^i(t)}{dt} = \overline{\left(\frac{\partial S(z)}{\partial z^i} \right)}.$$

$\langle \mathcal{K}_\sigma, \mathbb{R} \rangle$: intersection numbers of duals \mathcal{K}_σ and \mathbb{R}^n
 $(\mathcal{K}_\sigma = \{z(0) | z(\infty) = z_\sigma\})$.



[Witten, arXiv:1001.2933, 1009.6032]

Lefschetz thimble for Airy integral

Airy integral is given as

$$\text{Ai}(a) = \int_{\mathbb{R}} \frac{dx}{2\pi} \exp i \left(\frac{x^3}{3} + ax \right)$$

Complexify the integration variable: $z = x + iy$.



Integrand on \mathbb{R} , and on \mathcal{J}_1
($a = 1$)

Rewrite the Airy integral

There exists two Lefschetz thimbles \mathcal{J}_σ ($\sigma = 1, 2$) for the Airy integral:

$$\text{Ai}(a) = \sum_{\sigma} n_{\sigma} \int_{\mathcal{J}_{\sigma}} \frac{dz}{2\pi} \exp i \left(\frac{z^3}{3} + az \right).$$

n_{σ} : intersection number of the steepest ascent contour \mathcal{K}_{σ} and \mathbb{R} .





Figure: Lefschetz thimbles \mathcal{J} and duals \mathcal{K} ($a = 1e^{0.1i}, -1$)

Simplest model study to understand the sign problem

One-site Fermi Hubbard model

One-site Hubbard model:

$$\hat{H} = U\hat{n}_{\uparrow}\hat{n}_{\downarrow} - \mu(\hat{n}_{\uparrow} + \hat{n}_{\downarrow}).$$

Fock state gives the number density immediately:

$$\langle \hat{n} \rangle = \frac{1}{\beta} \frac{\partial}{\partial \mu} \ln Z = \frac{2(e^{\beta\mu} + e^{\beta(2\mu-U)})}{1 + 2e^{\beta\mu} + e^{\beta(2\mu-U)}}.$$

In the zero-temperature limit,

$$n(\beta = \infty) = \begin{cases} 2 & (1 < \mu/U), \\ 1 & (0 < \mu/U < 1), \\ 0 & (\mu/U < 0). \end{cases}$$

Path integral for one-site model

The path-integral expression for the one-site Hubbard model: :

$$Z = \sqrt{\frac{\beta}{2\pi U}} \int_{\mathbb{R}} d\varphi \underbrace{\left(1 + e^{\beta(i\varphi + \mu + U/2)}\right)^2}_{\text{Fermion Det}} e^{-\beta\varphi^2/2U}.$$

Integrand has complex phases causing the sign problem.

φ is an auxiliary field for the number density:

$$\langle \hat{n} \rangle = \text{Im} \langle \varphi \rangle / U.$$

Flows at $\mu/U < -0.5$ and $\mu/U > 1/5$




Figure: Flow at $\mu/U = 2$

$$Z = \int_{\mathcal{J}_*} dz e^{-S(z)}.$$

Number density: $n_* = 0$ for $\mu/U < -0.5$, $n_* = 2$ for $\mu/U > 1.5$.

(YT, Hidaka, Hayata, 1509.07146)

Flows at $-0.5 < \mu/U < 1.5$





Figure: Flow at $\mu/U = 0$

Complex saddle points lie on $\text{Im}(z_m)/U \simeq \mu/U + 1/2$.

This value is far away from $n = \text{Im} \langle z \rangle / U = 0, 1, \text{ or } 2$.

Curious incident of n in the one-site model

We have a big difference between the exact result and naive expectation:



(YT, Hidaka, Hayata, arXiv:1509.07146[hep-th])

(cf. Monte Carlo with one-thimble approx. gives the naively expected results.)

Fujii, Kamata, Kikukawa, 1509.08176, 1509.09141; Alexandru, Basar, Bedaque, 1510.03258)

Question

Can we unveil this curious incident?

Semiclassical study of the curious incident

Complex classical solutions

Let us solve, for $-0.5 < \mu/U < 1.5$,

$$\frac{\beta}{U} iz + \frac{2\beta \exp \beta \left(iz + \mu + \frac{U}{2} \right)}{1 + \exp \beta \left(iz + \mu + \frac{U}{2} \right)} = 0.$$

If $\beta U \gg 1$, the solutions are labeled by $m \in \mathbb{Z}$:

$$z_m \simeq i \left(\mu + \frac{U}{2} \right) + 2\pi m T.$$

At these solutions, the classical actions become

$$\begin{aligned} S_0 &\simeq -\frac{\beta U}{2} \left(\frac{\mu}{U} + \frac{1}{2} \right)^2, \\ \operatorname{Re}(S_m - S_0) &\simeq \frac{2\pi^2}{\beta U} m^2, \\ \operatorname{Im} S_m &\simeq 2\pi m \left(\frac{\mu}{U} + \frac{1}{2} \right). \end{aligned}$$

Semiclassical partition function

Using complex classical solutions z_m , let us calculate

$$Z_{\text{cl}} := \sum_{m=-\infty}^{\infty} e^{-S_m}.$$


This expression is valid for $-1/2 \lesssim \mu/U \lesssim 3/2$.

This is calculable using the elliptic theta function:

$$\begin{aligned} Z_{\text{cl}} &\simeq e^{-S_0} \left(1 + 2 \sum_{m=1}^{\infty} \cos 2\pi m \left(\frac{\mu}{U} + \frac{1}{2} \right) e^{-2\pi^2 m^2 / \beta U} \right) \\ &= e^{-S_0} \theta_3 \left(\pi \left(\frac{\mu}{U} + \frac{1}{2} \right), e^{-2\pi^2 / \beta U} \right). \end{aligned}$$

Number density & Lee–Yang zeros

Lee–Yang zeros of Z_{cl} :



Semiclassical study gives **the correct transition!**

$$n_{\text{cl}} := \frac{1}{\beta} \frac{\partial}{\partial \mu} \ln Z_{\text{cl}} \rightarrow \begin{cases} 2 & (1 < \mu/U < 3/2), \\ 1 & (0 < \mu/U < 1), \\ 0 & (-1/2 < \mu/U < 0). \end{cases}$$

(YT, Hidaka, Hayata, arXiv:1509.07146[hep-th])

Important interference among multiple thimbles

Let us consider a “phase-quenched” multi-thimble approximation:

$$Z_{|\text{cl.}|} = \sum_m |e^{-S_m}| = e^{-S_0(\mu)} \theta_3(0, e^{-2\pi^2/\beta U}).$$

- Lee–Yang zeros cannot appear at $\mu/U = 0, 1$.
- One-thimble, or “phase-quenched”, result: $n \simeq \mu/U + 1/2$.

Consequence


*In order to describe the step functions, we need **interference of complex phases** among different Lefschetz thimbles.*

(cf. Particle Productions: Dumulu, Dunne, PRL 104 250402)

(cf. Hidden Topological Angles: Behtash, Sulejmanpasic, Schäfer, Ünsal, PRL 115 041601)

Numerical results


Results for $\beta U = 30$: (1, 3, 5-thimble approx.: \mathcal{I}_0 , $\mathcal{I}_0 \cup \mathcal{I}_{\pm 1}$, and $\mathcal{I}_0 \cup \mathcal{I}_{\pm 1} \cup \mathcal{I}_{\pm 2}$)



(YT, Hidaka, Hayata, arXiv:1509.07146[hep-th])

Numerical results

Results for $\beta U = 100$:



Necessary numbers of Lefschetz thimbles $\propto \beta U / 2\pi$.

(YT, Hidaka, Hayata, arXiv:1509.07146[hep-th])

Summary and conclusion

Summary and Conclusion

- Picard–Lefschetz theory gives a suitable framework for saddle-point analysis even if $S(\phi)$ takes complex values.
- One-site Hubbard model is a nice toy model to play with the sign problem.
- Destructive interference among complex phases of Lefschetz thimbles play a pivotal role for the baryon Silver Blaze.